



Is the Betic-Rif Arc still collapsing?

J.I. Soto (1) and F. Fernandez-Ibañez (1)

(1) Instituto Andaluz de Ciencias de la Tierra & Departamento de Geodinámica, (CSIC-University of Granada), Campus Fuentenueva s/n, 18002-Granada, Spain (jsoto@ugr.es).

The Alpine orogenic arc formed by the Betics and Rif mountain ranges protects a close basin floored by an extremely attenuated continental crust: the Alboran Sea. It is demonstrated that westwards migration of the arc occurred simultaneously with extension in the Alboran region during the Neogene (mostly since the Oligocene and through the Miocene), within an oblique plate convergence between the African and Eurasian plates. Extensional collapse of the orogen was driven by low-angle extensional fault systems, which determined surface uplift and high denudation rates throughout the Miocene. Within this scenario we discuss if the relative motion of Africa and Eurasia plates controlled the extensional collapse of the Betic-Rif Arc, and if the collapse-related processes can be still operative in the region.

According to plate reconstructions in the Western Mediterranean region, we demonstrate that abrupt variations in the azimuth and magnitude of plate convergence during the Miocene conditioned the deformation in the arc, resulting that the main extensional collapse occurred immediately after an outstanding reduction of the Africa-Eurasia convergence rate.

Gathering information from diverse stress indicators (earthquake, borehole, and data from recent faults) we reconstruct the present-day stress field throughout the Gibraltar Arc. This field results from the complex superposition of two different stress sources; namely, differences in the gravitational potential energy imposed by the crustal structure and the occurrence of active faults. Extensional collapse in the arc is restricted currently to the central Betics, coinciding with the region with higher topographic elevation and maximum crustal thickness (>34 km). Besides this restricted domain, active deformation is accomplished by a major, left-lateral strike-slip fault system that cross cut entirely and obliquely the arc, from eastern Betics to the Rif. Stress rotations

along this fault zone and the overall stress field evidence that most of the plate convergence deformation is currently accommodated by lateral extrusion of material along this major-scale strike slip fault system, rather than ongoing extensional collapse of the Betic-Rif Arc.